

32-12099: Mouse Granulocyte Macrophage-Colony Stimulating Factor (AF)

Gene : Csf2
Gene ID : 12981
Uniprot ID : P01587
Alternative Name : Colony-stimulating factor, Csfgm

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 14.3 kDa (125 aa)

Granulocyte-macrophage colony-stimulating factor (GM-CSF) is hematopoietic growth factor produced by endothelial cells, monocytes, fibroblasts, and T cells. GM-CSF stimulates the production of neutrophilic granulocytes, macrophages, and mixed granulocyte-macrophage colonies from bone marrow cells. GM-CSF promotes immune system development and regulates neutrophil function during infection. Human and mouse GM-CSF show no cross-reactivity.

Product Info

Amount : 20 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at $\geq 95\%$
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM acetic acid Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MAPTRSPITV TRPWKHVEAI KEALNLLDDM PVTLNREEVEV VSNEFSFKKL TCVQTRLKIF EQGLRGNFTK LKGALNMTAS YYQTYCPPTP ETDCEQVTT YADFDISLKT FLTDIPFECK KPVQK

Application Note

Endotoxin: Less than $0.1 \text{ ng}/\mu\text{g}$ (1 IEU/ μg) as determined by LAL test.

Biological Activity was determined by FDCP-1 cell proliferation at $\leq 50 \text{ pg/mL}$; $\geq 2.0 \times 10^7 \text{ units/mg}$. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



