

32-12129: Mouse Interleukin-1 alpha (AF)

Gene : Il1a
Gene ID : 16175
Uniprot ID : P01582

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 18.1 kDa (157 aa)

Interleukin 1 alpha (IL-1alpha) is expressed by epithelial cells, activated macrophages, neutrophils, and endothelial cells to regulate immune responses. IL-1alpha signals through the IL-1 receptor, type 1 (IL-1R1) to activate the myeloid differentiation primary response 88 (MYD88) signaling pathway, which contains the cytoplasmic Toll/IL-1 receptor (TIR) domain adapter. IL-1alpha and the independently regulated IL-1beta protein have overlapping proinflammatory activities to induce adhesion molecule expression on epithelial cells, control fever induction, initiate rheumatoid arthritis, and promote septic shock.

Product Info

Amount : 20 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at $\geq 95\%$
Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5
Sterile water at 0.1 mg/mL
Content :
Storage condition : Store at -20°C
Amino Acid : MSAPYTYQSD LRYKLMKLV R QKFVMNDSL N QTIYQDV D K H YLSTTWLNDL QQEVKFD MYA YSSGGDDSKY PVTLKISDSQ L FVSAQGEDQ PVLLKELPET PKLITGSETD LIFFWKSINS KNYFTSAAYP ELFIATKEQS RVHLARGLPS MTD FQIS

Application Note

Endotoxin: Less than $0.1 \text{ ng}/\mu\text{g}$ ($1 \text{ IEU}/\mu\text{g}$) as determined by LAL test.

Biological Activity was determined by D10.G4.1 Proliferation. at $\leq 10 \text{ pg/mL}$; $\geq 1.0 \times 10^8 \text{ units/mg}$. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



