

32-12132: Mouse Interleukin-1 beta

Gene : Il1b
Gene ID : 16176
Uniprot ID : P10749
Alternative Name : Catabolin, LAF, EP, LEM, MCF

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 17.5 kDa (153 aa)

Interleukin 1 beta (IL-1beta) is a pro-inflammatory cytokine that is produced by monocytes, macrophages, and dendritic cells (DCs). IL-1beta signals through the IL-1 receptor, type 1 (IL-1R1) to activate the myeloid differentiation primary response 88 (MYD88) signaling pathway, which contains the cytoplasmic Toll/IL-1 receptor (TIR) domain adapter. IL-1beta and the independently regulated IL-1alpha protein have overlapping proinflammatory activities to induce adhesion molecule expression on epithelial cells, control fever induction, initiate rheumatoid arthritis, and promote septic shock.

Product Info

Amount : 10 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at >= 95%
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5
Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MVPIRQLHYR LRDEQKSLV LSDPYELKAL HLNQINQQ VIFSMSFVQG EPSNDKIPVA LGLKGNLYL SCVMKDGTP LQLESVDPKQ YPKKMEKRF VFNKIEVSK VEFESAEPN WYISTSQAEH KPVFLGNNSG QDIIDFTMES VSS

Application Note

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Biological Activity was determined by D10S cell proliferation at <=15 pg/mL; >= 6.7 x 10⁷ untis/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



