

32-12139: Human Interleukin-3

Gene : IL3
Gene ID : 3562
Uniprot ID : P08700

Alternative Name : Hematopoietic growth factor, Mast cell growth factor, Multipotential colony-stimulating factor, P-cell-stimulating factor

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 15.2 kDa (134 aa)

Interleukin 3 (IL-3) is a cytokine that is produced by activated T cells and mast cells. IL-3 induces the differentiation of hematopoietic stem cells into myeloid precursor cells, such as erythrocyte, megakaryocyte, granulocyte, monocyte, and dendritic cells. IL-3 also functions in the nervous system and is important during the B-1 cell regulation of chronic inflammatory diseases.

Product Info

Amount : 10 µg / 100 µg

Purification : Reducing and Non-Reducing SDS PAGE at $\geq 95\%$

Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA) or New Formulation with: Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.5 (available by request until we run out of the TFA formulation lots).

Sterile 10 mM acetic acid of 0.1 mg/mL for TFA formulation or Sterile water at 0.1 mg/mL for PBS formulation.

Storage condition : Store at -20°C

Amino Acid : MAPMTQTTPL KTSWVNCNSM IDEIITHLKQ PPLPLDFNN LNGEDQDILM ENNLRRPNLE AFNRAVKSLQ NASAIESILK NLLPCLPLAT AAPTRHPIHI KGDWNEFRR KLTFYKLTLE NAQAQQTLS LAIF

Application Note

Endotoxin: Less than $0.1 \text{ ng}/\mu\text{g}$ (1 IEU/ μg) as determined by LAL test.

Biological Activity was determined by TF-1 cell proliferation at $\leq 2 \text{ ng/mL}$; $\geq 5.0 \times 10^5 \text{ units/mg}$. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



