

36-1531: Monoclonal Antibody to PAX7 (Rhabdomyosarcoma Marker)(Clone : PAX7/1187)(Discontinued)

Clonality :	Monoclonal
Clone Name :	PAX7/1187
Application :	IHC
Reactivity :	Human
Gene :	PAX7
Gene ID :	5081
Uniprot ID :	P23759
Format :	Purified
Alternative Name :	PAX7,HUP1
Isotype :	Mouse IgG1 Kappa
Immunogen Information :	Recombinant fragment (aa300-600) of human PAX7 protein

Description


The Pax gene family of nuclear transcription factors is comprised of nine members that function during embryogenesis to regulate the temporal and position-dependent differentiation of cells. In addition, the family is involved in a variety of signal transduction pathways in the adult organism. Mutations in the Pax family of proteins have been linked to disease and cancer in humans. Pax-7 is a protein specifically expressed in cultured satellite cell-derived myoblasts. In situ hybridization reveals that Pax-7 is also expressed in satellite cells residing in adult muscle. A chromosomal aberration in the gene encoding Pax-7 causes rhabdomyosarcoma 2 (RMS2) (also called alveolar rhabdomyosarcoma).

Product Info


Amount :	100 µg
Purification :	Affinity Chromatography
Content :	100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
Storage condition :	Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid repeated freeze and thaw cycles.

Application Note

Immunohistology (Formalin-fixed) (0.5-1 µg/ml for 30 minutes at RT); (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris Buffer with 1mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes); Optimal dilution for a specific application should be determined.



Formalin-fixed, paraffin-embedded human Rhabdomyosarcoma stained with PAX7 Monoclonal Antibody (PAX7/1187).



Formalin-fixed, paraffin-embedded human Testicular Carcinoma stained with PAX7 Monoclonal Antibody (PAX7/1187).