

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-12176: Mouse Interleukin-13

Gene : II13 **Gene ID :** 16163 **Uniprot ID :** P20109

Alternative Name: T-cell activation protein P600, NC30

Description

Source: Genetically modified E.coli.

Predicted MW:Â Monomer, 12.3 kDa (111 aa)

Interleukin 13 (IL-13) is a cytokine secreted from type 2 T helper (Th2) cells. Â IL-13 has overlapping functions with interleukin 4 (IL-4), including the induction of immunoglobulin E (IgE) secretion from B cells, and the inhibition of interleukin 1 beta (IL-1beta), tumor necrosis factor alpha (TNF-alpha), interleukin 8 (IL-8), and interleukin 6 (IL-6) inflammatory cytokine expression. IL-13 also regulates immune cell inflammation in response to the pathophysiological changes of surrounding non-immune cells. The IL-13 receptor consists of the IL-4Ra and IL-13Ra1 subunits. IL-13 can also bind the IL-13Ra2 receptor with high affinity. IL-13 functions are mediated through the JAK/STAT signaling pathway. Human and mouse IL-13 are cross-reactive. Â

Product Info

Amount : $10 \mu g / 100 \mu g$

Purification: Reducing and Non-Reducing SDS PAGE at >= 95%

Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic

Content: Acid (TFA)

Sterile 20 mM HCl at 0.1 mg/mL

Storage condition : Store at -20°C


Amino Acid: MPVPRSVSLP LTLKELIEEL SNITQDQTPL CNGSMVWSVD LAAGGFCVAL DSLTNISNCN AIYRTQRILH

GLCNRKAPTT VSSLPDTKIE VAHFITKLLS YTKQLFRHGP F

Application Note

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.

Biological Activity was determined by TF-1 cell proliferation at <=10 ng/mL; $>=1.0 \times 10^5$ units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at $-80\text{\AA}\Box\text{\AA}^\circ\text{C}$ and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss.





9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

Mouse IL-13 Induced Proliferation of TF-1 Cells

1.20
1.00
0.80
0.80
0.40
0.20
0.00
0.01
0.10
1.00
1.000
100.00
100.00
100.00
100.00
100.00
100.00