

32-12177: Human Interleukin-15

Gene : IL15
Gene ID : 3600
Uniprot ID : P40933
Alternative Name : IL-15

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 12.9 kDa (115 aa)

Interleukin 15 (IL-15) is a widely expressed proinflammatory cytokine that is structurally and functionally related to interleukin 2 (IL-2). IL-15 signals through JAK kinases to activate STAT3, STAT5, and STAT6 transcription factors. IL-15 regulates the activation of T cells, neutrophils, and macrophages, and is a stimulatory cytokine promoting dendritic cell function. IL-15 expression is dysregulated in chronic inflammatory disorders, including rheumatoid arthritis, psoriasis, and pulmonary inflammatory diseases. Therefore, IL-15 may serve as an effective therapeutic target, due to the beneficial outcomes of IL-15 neutralization in models of psoriasis and diabetes. Human IL-15 shows activity on mouse cells.

Product Info

Amount : 10 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at $\geq 95\%$
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium bicarbonate, pH 8.0
Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MNWVNVISDL KKIEDLIQSM HIDATLYTES DVHPSCKVTA MKCFLLLELQV ISLESGDASI HDTVENLIL ANNSLSSNGN VTESGCKECE ELEEKNIKEF LQSFVHIVQM FINTS

Application Note

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Biological Activity was determined by CTLL-2 cell proliferation at ≤ 5 ng/mL; $\geq 2.0 \times 10^5$ units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



