

32-12184: Rat Interleukin-17A

Gene : Il17a
Uniprot ID : Q61453
Alternative Name : Ctl8, IL17, Cytotoxic T-lymphocyte-associated antigen 8

Description

Source: Genetically modified E.coli.

Predicted MW: Dimer, 15.1/30.3 kDa (134/268 aa)

IL-17A (IL-17A), also known as CTLA-8, is a member of the IL-17 family of proteins. IL-17A is proinflammatory cytokine that is secreted by activated CD4+ and CD8+ T lymphocytes. IL-17A acts through its receptor, IL-17R, to promote increased cytokine and chemokine secretion. In turn, the cytokines and chemokines mediate the immunoregulatory function of IL-17A by promoting the proliferation, maturation, and chemoattraction of neutrophils to inflammatory sites. Elevated levels of IL-17A are associated with rheumatoid arthritis, airway inflammation, allograft rejection, inflammatory bowel disease, psoriasis, cancer, and multiple sclerosis. Human, mouse, and rat IL-17A show activity on mouse cells.

Product Info

Amount : 25 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at >= 95%
Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium citrate, pH 3.0
Content : Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MAVLIPQSSV CPNAEANNFL QNVKVNKVL NSLSSKASSR RPSDYLNRSST SPWTLSRNEP PDRYPSVIWE AQRHRQRCVN AEGKLDHMM SVLIQQEILV LKREPEKCPF TFRVEKMLVG VGCTCVSSIV RHAS

Application Note

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



