

## 32-12189: Mouse Interleukin-17E

**Gene :** Il25  
**Uniprot ID :** Q8VHC9  
**Alternative Name :** Il17e, Interleukin-17E

### Description

**Source:** Genetically modified E.coli.

**Predicted MW:** Dimer, 17.6/35.2 kDa (154/308 aa)

IL-17E (IL-17E), also known as IL-25, is a member of the IL-17 family of cytokines. IL-17E binds to the IL-17RB receptor to stimulate the secretion of the proinflammatory interleukin 8 (IL-8) chemokine and to induce the activation of nuclear factor kappa-light-chain-enhancer of activated B cells (NF- $\kappa$ B).

### Product Info

**Amount :** 25  $\mu$ g / 100  $\mu$ g  
**Purification :** Reducing and Non-Reducing SDS PAGE at  $\geq$  90%  
Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)  
**Content :** Sterile 10 mM HCl at 0.1 mg/mL  
**Storage condition :** Store at -20°C  
**Amino Acid :** MVSLRIQEGC SHLPSCCPSK EQEPPEEWLK WSSASVSPPE PLSHTTHAES CRASKDGPLN SRAISPWSYE LDRDLNRVPQ DLYHARCLCP HCVSLQTGSH MDPLGNSVPL YHNQTVFYRR PCHGEEGTHR RYCLERRLYR VSLACVCRP RVMA

### Application Note

**Endotoxin:** Less than 0.1 ng/ $\mu$ g (1 IEU/ $\mu$ g) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.

