# **w** abeomics

#### 32-12201: Human Interleukin-21

Gene :IL21Gene ID :59067Uniprot ID :Q9HBE4Alternative Name :Za11, Interleukin-21Â Â

### Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 15.4 kDa (132 aa)

Interleukin 21 (IL-21) is a member of the common-gamma chain family of cytokines that are involved in immunoregulation. IL-21 is normally expressed by activated CD4+ T cells and is aberrantly expressed in Hodgkin lymphoma cells. The IL-21 receptor (IL-21R) activates the JAK/STAT signaling pathway and is expressed on T, B, and natural killer (NK) cells. Within the B cell lineage, IL-21 is a switch factor regulating IgG1 and IgG3 antibody production. IL-21 also cooperates with IL-4 for the production of multiple antibody classes in B cells. IL-21 has pleiotropic effects on the proliferation, differentiation, and effector functions of B, T, NK, and dendritic cells.


### **Product Info**

Amount :	10 μg / 100 μg
Purification :	Reducing and Non-Reducing SDS PAGE at $>= 95\%$
Content :	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5 Sterile water at 0.1 mg/mL
Storage condition :	Store at -20°C
Amino Acid :	MQDRHMIRMR QLIDIVDQLK NYVNDLVPEF LPAPEDVETN CEWSAFSCFQ KAQLKSANTG NNERIINVSI KKLKRKPPST NAGRRQKHRL TCPSCDSYEK KPPKEFLERF KSLLQKMIHQ HLSSRTHGSE DS

### **Application Note**


**Endotoxin:** Less than 0.1 ng/ $\mu$ g (1 IEU/ $\mu$ g) as determined by LAL test.

Biological Activity was determined by ANBL-6 cell proliferation OR B9 cell proliferation at ED50 <= 2 ng/mL; >=  $5.0 \times 10^{5}$  units/mg (typical ED50 is < 1 ng/mL) OR ED50 <= 50 ng/mL; >=  $2.0 \times 10^{4} \text{ units/mg}$ 



# **₩** abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



For Research Use Only. Not for use in diagnostic/therapeutics procedures.