

32-12203: Mouse Interleukin-21

Gene : IL21
Gene ID : 60505
Uniprot ID : Q9ES17
Alternative Name : ⌘ ⌘ ⌘ ⌘ ⌘

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 15.1 kDa (130 aa)

Interleukin 21 (IL-21) is a member of the common-gamma chain family of cytokines that are involved in immunoregulation. IL-21 is normally expressed by activated CD4+ T cells. The IL-21 receptor (IL-21R) activates the JAK/STAT signaling pathway and is expressed on T, B, and natural killer (NK) cells. Within the B cell lineage, IL-21 is a switch factor regulating IgG1 and IgG3 antibody production. IL-21 also cooperates with interleukin 4 (IL-4) for the production of multiple antibody classes in B cells. IL-21 has pleiotropic effects on the proliferation, differentiation, and effector functions of B, T, NK, and dendritic cells.

Product Info

Amount : 10 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at >= 95%
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 20 mM sodium bicarbonate, pH 8.5
Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MHKSSPQGPD RLLIRLRHLI DIVEQLKIYE NDLDPSELLSA PQDVKGHCEH AAFACFQKAK LKPSNPGNNK TFIIDLVAQL RRRLPARRGG KKQKHIKCP SCDSYEKTRP KEFLERLKWL LQKMIHQHLS

Application Note

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Biological Activity was determined by Mouse thymocyte proliferation with costimulation at <=10 ng/mL; >= 1.0 x 10⁵ units/mg . Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



