

## 32-12215: Human IP-10 (CXCL10)

**Gene :** CXCL10

**Gene ID :** 3627

**Uniprot ID :** P02778

**Alternative Name :** C-X-C motif chemokine 10, 10 kDa interferon gamma-induced protein, Small-inducible cytokine B10

### Description

**Source:** Genetically modified E.coli.

**Predicted MW:** Monomer, 8.6 kDa (77 aa)

Interferon gamma-induced protein 10 (IP-10), or CXCL10, is a chemokine secreted by monocytes, endothelial cells and fibroblasts in response to interferon gamma (IFN-gamma). IP-10 functions as a chemoattractant for activated T cells, monocytes, dendritic, and natural killer (NK) cells that express the G protein-coupled receptor CXCR3. IP-10 is an important factor in autoimmune diseases such as Hashimoto's thyroiditis, Graves' disease, and Type 1 diabetes mellitus.

### Product Info

**Amount :** 25 µg / 100 µg

**Purification :** Reducing and Non-Reducing SDS PAGE at >= 95%

**Content :** Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)  
Sterile water at 0.1 mg/mL

**Storage condition :** Store at -20°C

**Amino Acid :** VPLSRTVRCT CISISNPVN PRSLEKLEII PASQFCPRVE IATMKKKGE KRCLNPESKA IKNLLKAVSK  
ERSKRSP

### Application Note

**Endotoxin:** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



