

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-12240: Mouse Macrophage-Colony Stimulating Factor (AF)

 Gene :
 Csf1

 Gene ID :
 12977

 Uniprot ID :
 P07141

Alternative Name: Macrophage colony-stimulating factor 1, CSF-1Â Â Â Â Â Â Â Â Â Â Â Â Â Â

Description

Source: Genetically modified E.coli.

Predicted MW: Dimer, 18.2/36.4 kDa (156/312 aa)

Macrophage colony stimulating factor (M-CSF) is a hematopoietic growth factor that is widely produced by a variety of cells. Â M-CSF stimulates the proliferation and differentiation of hematopoietic stem cells into monocyte and macrophage cell types. Â M-CSF also acts through the colony stimulating factor 1 receptor (CSF1R) to modulate processes involved in immunology, bone metabolism, fertility, and pregnancy. Â Human M-CSF shows activity on mouse cells; however, mouse M-CSF shows no activity on human cells. Â

Product Info

Amount : $10 \mu g / 100 \mu g$

Purification: Reducing and Non-Reducing SDS PAGE at >= 95%

Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium

Content: phosphate, 50 mM sodium chloride, pH 7.5

Sterile water at 0.1 mg/mL

Storage condition : Store at -20°C

Amino Acid: MKEVSEHCSH MIGNGHLKVL QQLIDSQMET SCQIAFEFVD QEQLDDPVCY LKKAFFLVQD IIDETMRFKD


NTPNANATER LQELSNNLNS CFTKDYEEQN KACVRTFHET PLQLLEKIKN FFNETKNLLE KDWNIFTKNC

NNSFAKCSSR DVVTKP

Application Note

Endotoxin: Less than 0.1 ng/ \tilde{A} $\square \hat{A}\mu g$ (1 IEU/ \tilde{A} $\square \hat{A}\mu g$) as determined by LAL test.

Biological Activity was determined by NFS-60 cell proliferation at <=10 ng/mL; $>=1.0 \times 10^5$ units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80\AA \square Å°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss.





9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

Mouse M-CSF Induced Proliferation of M-NFS-60 Cells

1.75
1.55
1.35
0.05
0.75
0.05
0.00
0.00
0.01
0.10
1.00
10.00
10.00
10.00