

32-12294: Mouse RELM-alpha

Gene : Retnla

Gene ID : 57262

Uniprot ID : Q9EP95

Alternative Name : Cysteine-rich secreted protein A12-gamma, I Cysteine-rich secreted protein FIZZ1, Hypoxia-induced mitogenic factor, Parasite-induced macrophage novel gene 1 protein, RELMalpha

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 9.6 kDa (89 aa)

Resistin-like molecule-alpha (RELM-alpha) is a member of the RELM family of secreted proteins containing conserved C-terminus cysteines. The RELM family consists of Resistin (FIZZ3), RELM-alpha (FIZZ1), RELM-beta (FIZZ2), and RELM-gamma (FIZZ4). RELM-alpha and Resistin are secreted by adipocytes, unlike RELM-beta which is secreted by gastrointestinal epithelial cells, and RELM-gamma which is expressed by peripheral blood granulocytes and bone marrow cells.

Product Info

Amount : 25 µg / 100 µg

Purification : Reducing and Non-Reducing SDS PAGE at >= 95%

Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5
Sterile water at 0.1 mg/mL

Storage condition : Store at -20°C

Amino Acid : MDETIEIIVE NKVKELLANP ANYPSTVTKT LSCTSVKTMN RWASCPAGMT ATGCACGFAC GSWEIQSGDT CNCLLLVDW TTARCCQLS

Application Note

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.

