

## 32-12303: Human Stem Cell Factor (AF)

**Gene :** KITLG  
**Gene ID :** 4254  
**Uniprot ID :** P21583  
**Alternative Name :** Resistin

### Description

**Source:** Genetically modified E.coli.

**Predicted MW:** Monomer, 18.6 kDa (165 aa)

Stem cell factor (SCF) is a cytokine made by fibroblasts and endothelial cells. SCF binds to the receptor c-Kit/CD117 and plays a critical role in the maintenance, survival, and differentiation of hematopoietic stem cells. While human SCF shows no activity on murine cells, murine and rat SCF are active on human cells.

### Product Info

**Amount :** 10 µg / 100 µg  
**Purification :** Reducing and Non-Reducing SDS PAGE at  $\geq 95\%$   
Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM acetic acid or New Formulation with Lot 0417-100-03 and 0417-100-03AF and moving forward: Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5 (available by request until we run out of acetic acid formulation lots).  
**Content :** Sterile 10 mM HCl at 0.1 mg/mL for acetic acid formulation or Sterile water at 0.1 mg/mL for PBS formulation.  
**Storage condition :** Store at  $-20^{\circ}\text{C}$   
**Amino Acid :** MEGICRNRVT NNVKDVTKLV ANLPKDYMIT LKYVPGMDVL PSHCWISEMV VQLSDSLTDL LDKFSNISEG LSNYSIIDKL VNI VDDLVEC VKENSSKDLK KSKFSPEPRL FTPEEFFRIF NRSIDAFKDF VVASETSDCV VSSTLSPEKD SRVSVTKPFM LPPVA

### Application Note

**Endotoxin:** Less than  $0.1 \text{ ng}/\mu\text{g}$  (1 IEU/ $\mu\text{g}$ ) as determined by LAL test.

Biological Activity was determined by TF-1 cell proliferation at  $\leq 15 \text{ ng/mL}$ ;  $\geq 6.6 \times 10^4 \text{ units/mg}$ . Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at  $-80^{\circ}\text{C}$  and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



