

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-12303: Human Stem Cell Factor (AF)

 Gene :
 KITLG

 Gene ID :
 4254

 Uniprot ID :
 P21583

Alternative Name: Resistin Â Â Â

## **Description**

Source: Genetically modified E.coli.

Predicted MW:Â Monomer, 18.6 kDa (165 aa)

Stem cell factor (SCF) is a cytokine made by fibroblasts and endothelial cells. SCF binds to the receptor c-Kit/CD117 and plays a critical role in the maintenance, survival, and differentiation of hematopoietic stem cells. While human SCF shows no activity on murine cells, murine and rat SCF are active on human cells.

## **Product Info**

**Amount :**  $10 \mu g / 100 \mu g$ 

**Purification:** Reducing and Non-Reducing SDS PAGE at >= 95%

Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM acetic acid or New Formulation with Lot 0417-100-03 and 0417-100-03AF and moving forward: Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, 50 mM sodium shloride and 7.5 (available by request until two run out of acetic acid formulation)

**Content :** mM sodium chloride, pH 7.5 (available by request until we run out of acetic acid formulation

lots).

Sterile 10 mM HCl at 0.1 mg/mL for acetic acid formulaiton or Sterile water at 0.1 mg/mL for

PBS formulation.

**Storage condition :** Store at -20°C

Amino Acid: MEGICRNRVT NNVKDVTKLV ANLPKDYMIT LKYVPGMDVL PSHCWISEMV VQLSDSLTDL LDKFSNISEG


LSNYSIIDKL VNIVDDLVEC VKENSSKDLK KSFKSPEPRL FTPEEFFRIF NRSIDAFKDF VVASETSDCV

VSSTLSPEKD SRVSVTKPFM LPPVA

## **Application Note**

**Endotoxin:** Less than 0.1 ng/ $\tilde{A}$  $\square \hat{A}\mu g$  (1 IEU/ $\tilde{A}$  $\square \hat{A}\mu g$ ) as determined by LAL test.

Biological Activity was determined by TF-1 cell proliferation at <=15 ng/mL; >= 6.6 x 10^4 units/mg  $\tilde{\mathbb{A}}$  . Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80 $\tilde{\mathbb{A}}$  $\tilde{\mathbb{A}}$ °C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss.





9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

