

32-12304: Mouse Stem Cell Factor

Gene : Kitlg
Gene ID : 17311
Uniprot ID : P20826
Alternative Name : Kit ligand, Hematopoietic growth factor KL, Mast cell growth factor, Steel factor, Stem cell factor, c-Kit ligand

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 18.4 kDa (165 aa)

Stem cell factor (SCF) is a cytokine made by fibroblasts and endothelial cells. SCF binds to the receptor c-Kit/CD117 and plays a critical role in the maintenance, survival, and differentiation of hematopoietic stem cells. While human SCF shows no activity on murine cells, murine and rat SCF are active on human cells.

Product Info

Amount : 10 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at >= 95%
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
 Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MKEICGNPVT DNVKDITKLV ANLPNDYMIT LNYVAGMDVL PSHCWLRDMV IQLSLSLTTL LDKFSNISEG
 LSNYSIIDKL GKIVDDLVLVLC MEENAPKNIK ESPKRPETRS FTPEEFSIF NRSIDAFKDF MVASDTSDCV
 LSSTLGPEKD SRVSVTKPFM LPPVA

Application Note

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Biological Activity was determined by TF-1 cell proliferation at <= 20 ng/mL; >= 5.0 x 10⁴ units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



