

## 32-12314: Human TARC (CCL17)

**Gene :** CCL17

**Gene ID :** 6361

**Uniprot ID :** Q92583

**Alternative Name :** C-C motif chemokine 17, C-C chemokine TARC, Small-inducible cytokine A17, Thymus and activation-regulated chemokine

### Description

**Source:** Genetically modified E.coli.

**Predicted MW:** Monomer, 8.1 kDa (71 aa)

Thymus and activation regulated chemokine (TARC), also known as CCL17, is a chemokine that is constitutively produced by thymus tissue and activated peripheral blood mononuclear cells (PBMCs), including dendritic cells. TARC signals through the CCR4 receptor to induce chemotaxis of Type 2 T helper (Th2) cells. TARC is important in asthma and allergic diseases, along with bacterial and viral infections.

### Product Info

**Amount :** 20 µg / 100 µg

**Purification :** Reducing and Non-Reducing SDS PAGE at >= 95%

**Content :** Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)  
Sterile water at 0.1 mg/mL

**Storage condition :** Store at -20°C

**Amino Acid :** ARGTVNGREC CLEYFKGAIP LRKLKTWYQT SEDCSRDAIV FVTVQGRAIC SDPNNKRVKV AVKYLQSLER S

### Application Note

**Endotoxin:** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.

