

32-12315: Human Transforming Growth Factor-beta 1

Gene : TGFB1
Gene ID : 7040
Uniprot ID : P01137
Alternative Name : Transforming growth factor beta-1, TGF-beta-1

Description

Source: Genetically modified HEK 293 cells.

Å Dimer, 12.8/25.6 kDa (112/224 aa)

Transforming growth factor-beta 1 (TGF-b1) is a member of the TGF-beta superfamily of cytokines.Å Members of this family exhibit regulatory activity in immunity, proliferation, adhesion, migration, and growth inhibition pathways.Å TGF-b1 signals through SMAD proteins via the TGF-bRI and TGF-bRII receptors.Å

Product Info

Amount : 20 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at >= 95%
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1 % trifluoroacetic acid (TFA) and Trehalose in a 20:1 Trehalose to protein ratio
Sterile 10 mM HCl at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : ALDTNYCFSS TEKNCCVRQL YIDFRKDLGW KWIHEPKGYH ANFCLGPCPY IWSLDTQYSK VLALYNQHNP GASAAPCCVP QALEPLPIVY YVGRKPKVEQ LSNMIVRSCK CS

Application Note

Endotoxin: Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.

Biological Activity was determined by Inhibition of IL-4-induced HT-2 cell proliferation at <=500 pg/mL; >= 2.0 x 10⁶ units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80Å°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialled to compensate for this loss.



