

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-12321: Human Tumor Necrosis Factor-alpha

 Gene :
 TNF

 Gene ID :
 7124

 Uniprot ID :
 P01375

Alternative Name: Tumor necrosis facto, Cachectin, TNF-alpha, Tumor necrosis factor ligand superfamily member 2,

NTFÂ Â Â Â Â Â

## **Description**

Source: Genetically modified E.coli.

Predicted MW: Monomer, 17.5 kDa (158 aa)

Tumor necrosis factor alpha (TNF-alpha) is an inflammatory cytokine secreted by macrophages, monocytes, neutrophils, T cells, and NK-cells following stimulation by bacterial lipopolysaccharide (LPS). TNF-alpha signal activation occurs through two receptors, TNFR1 and TNFR2. TNFR1 is expressed on most cell types, unlike TNFR2, which is expressed mainly on immune cells. TNF-alpha functions to stimulate phagocytosis in macrophages, chemoattract neutrophils, increase insulin resistance, and induce fever.

## **Product Info**

**Amount:** 50 μg / 100 μg

**Purification:** Reducing and Non-Reducing SDS PAGE at >= 95%

Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium

**Content:** phosphate, pH 7.5

Sterile water at 0.1 mg/mL

**Storage condition :** Store at -20°C

Amino Acid: MVRSSSRTPS DKPVAHVVAN PQAEGQLQWL NRRANALLAN GVELRDNQLV VPSEGLYLIY

SOVLFKGOGC PSTHVLLTHT ISRIAVSYOT KVNLLSAIKS PCORETPEGA EAKPWYEPIY LGGVFOLEKG

DRLSAEINRP DYLDFAESGQ VYFGIIAL

## **Application Note**


Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Biological Activity was determined by Cytolysis of mouse L929 cells in the presence of Actinomycin D at <=2 ng/mL; >=5.0 x  $10^5$  units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at  $-80^\circ$ C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



Human TNF alpha
Figure: 1 ug in each lane (-) non-reducing conditions and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Human TNF alpha has a predicted MW of 17.5 kDa.

