

## 10-3014: Monoclonal antibody to TLR5 (Clone: ABM22C2)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	ABM22C2
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Gene :</b>	TLR5
<b>Gene ID :</b>	7100
<b>Uniprot ID :</b>	O60602
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Toll/interleukin-1 receptor-like protein 3, TIL3
<b>Isotype :</b>	Mouse IgG2a, Kappa
<b>Immunogen Information :</b>	A partial length recombinant TLR5 protein was used as the immunogen for this antibody.

### Description

Toll-like receptor 5 (TLR5) plays a fundamental role in pathogen recognition and downstream signaling to induce effective immunity. It is well expressed in major old tissues including lung, skin, intestine, and spleen. TLR5 is present in synovial of patients with rheumatoid arthritis (RA), act as novel mediator of innate immunity-induced osteoclastogenesis and bone loss. TLR5 is a receptor for flagellin and is present on the basolateral surface of intestinal epithelial cells. It is involved in the migration of intestinal epithelial cells through activation of the p38 MAPK pathway. TLR5-mediated signaling specifically enhanced the maturation of lung dendritic cells. TLR5 influences the progression and outcome of ovarian, sarcoma, and luminal breast tumors. Mutations in TLR5 have been associated with both resistance and susceptibility to systemic lupus erythematosus, and susceptibility to Legionnaire disease.

### Product Info

<b>Amount :</b>	25 µg / 100 µg
<b>Purification :</b>	Protein G Chromatography
<b>Content :</b>	25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.
<b>Storage condition :</b>	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.

### Application Note

FlowCytometry: 0.5-1 µg/10<sup>6</sup> Cells




Figure-1: Cell surface flowcytometry analysis of hTLR5 on human PBMC (Monocytes) using 0.5  $\mu\text{g}/10^6$  Cells of Anti-TLR5 antibody (Clone: ABM22C2). Green represent isotype control and red represent Anti human TLR5 antibody (Abeomics 10-3014). Goat anti mouse PE conjugated was used as the secondary antibody.