

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

45-1015: Rabbit Polyclonal Antibody to CBP-tag

Clonality: Polyclonal
Application: ELISA, WB
Format: Purified
Isotype: Rabbit IgG

Immunogen Information: A synthetic CBP tag peptide KRRWKKNFIAVSAANRFKKISSSGAL conjugated to KLH

Description

CBP tag is a calmodulin-binding peptide, KRRWKKNFIAVSAANRFKKISSSGAL, derived from skeletal/cardiac muscle myosin light chain kinase 2. Rabbit Anti-CBP-tag Polyclonal Antibody is developed in rabbit hosts using a synthetic CBP tag peptide conjugated to KLH. CBP-tag Antibody is highly purified from rabbit antiserum by immunoaffinity chromatography. The antibody is suitable for detecting CBP fusion proteins.

Product Info

Amount: $40 \mu g$

Purification: Immunoaffinity chromatography

Content: 0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide

The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody

Storage condition : can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or

below. Avoid repeated freezing and thawing cycles.

Application Note

ELISA: 0.05-0.2 ̸Âμg/ml Western blot: 1-2 ̸Âμg/ml

Reconstitute the lyophilized powder with deionized water (or equivalent) to an antibody concentration of 0.5

mg/ml.




Figure-1 : Western blot analysis of CBP-tag Antibody at 1 μ g/ml on CBP tagged fusion protein (200 & 40 ng) expressed in E. coli Cell Lysate.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-2 : Western blot analysis of CBP-tag Antibody at 1 μ g/ml on CBP tagged fusion protein (50, 20, 10 & 5 ng)expressed in E. coli cell lysate. IRDy 800 Conjugated Goat Anti-Rabbit IgG was used as Secondary Antibody.




Figure-3: Western blot analysis of CBP-tag Antibody on CBP-tagged protein, Lane-1: CBP-tagged protein from the transfected CHO cell lysates (as input control), Lane-2: Immuno-precipitates of CBP-tagged protein from the transfected CHO cell lysates with CBP Tag Antibody, 3: Immuno-precipitates of the CBP-tagged protein from the transfected CHO cell lysates with Rabbit IgG Control (Whole Molecule).