w abeomics

45-1019: Rabbit Polyclonal Antibody to DDDDK-tag (Discontinued)

Clonality :	Polyclonal
Application :	ELISA
Format :	Purified
Isotype :	Rabbit IgG
Immunogen Information :	A synthetic epitope peptide xxxDDDDK conjugated to KLH. This sequence represents the enterokinase cleavage site .

Description

Well-characterized antibodies for epitope tags consisting of short sequences have been widely used in the study of protein expression in various systems. The xxxDDDDK-tag is a small size and high hydrophilicity epitope tag containing an enterokinase cleavage site that allows it to be completely removed from the fusion proteins. The Rabbit Anti-DDDDK-tag Polyclonal Antibody is developed in rabbit hosts using a synthetic peptide containing xxxDDDDK epitope conjugated to KLH. This polyclonal antibody is highly purified from rabbit antiserum by immunoaffinity chromatography.

Product Info

Amount :	40 µg
Purification :	Immunoaffinity chromatography
Content :	0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide.
Storage condition :	The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or below. Avoid repeated freezing and thawing cycles.

Application Note

ELISA: 0.05-0.2 μg/ml **Western blot:**1-2 μg/ml



Figure-1 : Western blot analysis of DDDDK-tag Antibody on DDDDK tagged fusion protein in Lane 1-3 (10, 1 & 0.1 ng) expressed in E. coli lysates, Lane- 4: E. coli lysates as Negative Control.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



Figure-2 : Western blot analysis of DDDDK-tag Antibody at 1 μ g/ml on DDDDK tagged fusion protein in Lane 1-3 (10, 1 & 0.1 ng) expressed in E. coli lysates, Lane-4: E. coli lysates as Negative Control.