

45-1071: Mouse Monoclonal Antibody to Phosphotyrosine (Clone : 18E10D2)(Discontinued)

Clonality :	Monoclonal
Clone Name :	18E10D2
Application :	WB
Reactivity :	Human
Format :	Purified
Isotype :	Mouse IgG2a, Kappa
Immunogen Information :	Phosphotyrosine containing carrier protein

Description

Phosphotyrosine is a tyrosine residue covalently bound to a phosphate via its hydroxyl group. A tyrosine kinase catalyzes the transfer of a phosphate from ATP to a tyrosine residue on the protein substrate. Tyrosine phosphorylation plays a key role in intracellular and cancer development. Many tyrosine kinases are drug targets for different forms of cancer. Phosphotyrosine antibody specifically binds to phosphotyrosine residues. It is a valuable tool to analyze tyrosine phosphorylation and monitor the activity of tyrosine kinase in high throughput drug discovery. Phosphotyrosine Antibody (18E10), mAb, Mouse is produced from a hybridoma resulting from the fusion of SP2/0-Ag14 myeloma and B-lymphocytes harvested from mouse immunized with phosphotyrosine containing carrier protein.

Product Info

Amount :	100 µg
Purification :	Protein A chromatography
Content :	0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide.
Storage condition :	The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or below. Avoid repeated freezing and thawing cycles.

Application Note

Western Blot: 1-2 µg/ml.




Figure-1 : Western blot analysis of Phosphotyrosine Antibody (Clone: 18E10D2) at 1 µg/ml on EGF-stimulated A431 cell lysates and untreated A431 cell lysates, IRDye 800 Conjugated affinity Purified Goat Anti-Mouse IgG was used as secondary Antibody.