

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 45-1079: Mouse Monoclonal Antibody to Human TIM-3 (Clone: TI.B2) (Discontinued)

Clonality: Monoclonal

Clone Name: TI.B2
Application: ELISA
Gene: HAVCR2
Gene ID: 84868
Uniprot ID: Q8TDQ0
Format: Purified

Alternative Name: T-Cell Immunoglobulin Mucin Receptor 3/HAVCR2/ CD366/ KIM-3/ TIMD3/ Tim-3/ TIMD-3/

HAVcr-2

**Isotype:** Mouse IgG1, Kappa

Immunogen Information: Recombinant human TIM3-Fc

## **Description**

TIM-3 (T cell immunoglobulin and mucin domain-3), is a member of the TIM family of immune regulating molecules. Its binding to Galectin-9 induces a range of immunosuppressive functions which enhance immune tolerance and inhibit antitumor immunity. Anti-Human TIM3 Antibody (TI.B2), mAb, Mouse is produced from the hybridoma resulting from fusion of SP2/0 myeloma and B-lymphocytes obtained from mouse immunized with recombinant human TIM3-Fc.

## **Product Info**

**Amount :** 40 μg

**Purification:** Protein A chromatography

**Content:** 0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide.

The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody

**Storage condition:** can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or

below. Avoid repeated freezing and thawing cycles.

## **Application Note**

ELISA detection:  $0.01\text{-}0.1 \,\mu\text{g/ml}$ Flow cytometry: 5-7  $\,\mu\text{g/ml}$ 




Figure-1 : ELISA binding of TIM3 Antibody (Clone: TI.B2) with Human TIM3 recombinant protein. Coating antigen: TIM3-Fc at 0.5  $\mu$ g/ml. TIM3 antibody dilution start from 1000 ng/ml, EC50= 6.98 ng/ml



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-2 : Flow cytometric analysis of TIM3 Antibody (Clone: TI.B2) on CHO-K1/TIM3 stable cell expressing TIM3 (Red histogram) and CHO negative control cell (Black histogram) at 5  $\mu$ g/ml, 2.5x10â $\mu$ cells/reaction, iFluor647 conjugated Goat Anti-Mouse IgG used as Secondary Antibody.