

45-1093: Mouse Monoclonal Antibody to Human OX40 (Clone : OX.A10) (Discontinued)

Clonality :	Monoclonal
Clone Name :	OX.A10
Application :	ELISA
Format :	Purified
Isotype :	Mouse IgG2b,Kappa
Immunogen Information :	Recombinant human OX40-His

Description

OX40 (CD134; TNFRSF4) is a T cell co-stimulatory molecule of the TNF receptor superfamily that coordinates with other co-stimulators (CD28, CD40, CD30, CD27 and 4-1BB) to manage the activation of the immune response. OX40 is up-regulated on CD4+ and CD8+ T cells upon engagement of the TCR by antigen presenting cells along with co-stimulation by CD40-CD40 Ligand and CD28-B7. OX40 Ligand is primarily expressed on antigen presenting cells. OX40 Ligand engagement of OX40 on activated CD4+ T cells results in increased T cell survival, proliferation, and cytokine production. It also inhibits the conversion of effector T cells into immunosuppressive regulatory T cells (Tregs) and can promote the maintenance of and recall response in memory T cells. OX40 is constitutively expressed on Tregs and enhances the sensitivity of Tregs to IL-2, thus promoting Treg proliferation. OX40 has also been shown to decrease the cells' immunosuppressive activity on effector T cells. OX40-OX40 Ligand signaling is involved in allergic airway inflammation, graft-versus-host disease and autoimmune disease. Mutations in OX40 and OX40 Ligand are associated with cardiovascular disease. Human OX40 Antibody (OX.A10), mAb, Mouse is produced from a hybridoma resulting from the fusion of SP2/0 myeloma and B-lymphocytes obtained from a mouse immunized with recombinant human OX40-His.

Product Info

Amount :	40 µg
Purification :	Protein A chromatography
Content :	0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide.
Storage condition :	The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or below. Avoid repeated freezing and thawing cycles.

Application Note

ELISA detection: 0.01-0.1 µg/ml

ELISA blocking: 10-15 µg/ml




Figure-1 : ELISA binding of OX40 Antibody (Clone: OX.A10) with Human OX40 recombinant protein, Coating antigen: OX40-His at 0.5 μ g/ml, Human OX40 Antibody (OX.A10) dilution start from 1000 ng/ml, EC50= 4.957 ng/ml.