

## 45-1095: Rabbit Monoclonal Antibody to DYKDDDDK Tag (Clone : 114F12C8)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	114F12C8
<b>Application :</b>	ELISA
<b>Format :</b>	Purified
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	DYKDDDDK synthetic peptide coupled to KLH

### Description

The rabbit immune system generates antibody diversity and optimizes affinity. technology to generate the high affinity and specificity monoclonal rabbit antibodies. DYKDDDDK Tag Antibody, mAb, Rabbit specific to DYKDDDDK tags placed at C-terminal, N-terminal and internal regions of fusion proteins. The antibody can greatly improve the effectiveness of several different kinds of immunoassays, helping researchers identify, detect, and purify DYKDDDDK fusion proteins in bacteria and mammalian cells.

### Product Info

<b>Amount :</b>	40 µg
<b>Purification :</b>	Protein A chromatography
<b>Content :</b>	0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide.
<b>Storage condition :</b>	The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or below. Avoid repeated freezing and thawing cycles.

### Application Note

ELISA: 0.05-0.2 µg/ml  
Western Blot: 0.2-1 µg/ml




Figure-1 : ELISA binding of DYKDDDDK Tag Antibody (Clone: 114F12C8) on DYKDDDDK Tagged protein, Coating antigen: DYKDDDDK-tagged fusion protein at 1 µg/ml, DYKDDDDK Tag Antibody dilution start from 1,000 ng/ml, EC50= 47.78 ng/ml.




Figure-2 : Western blot analysis of DYKDDDDK Tag Antibody (Clone: 114F12C8) at 0.2  $\mu$ g/ml on N-terminal of DYKDDDDK-tagged fusion protein (1-4: 25 ng, 10 ng, 5 ng & 1 ng respectively), C-terminal of DYKDDDDK-tagged fusion protein (5-8: 25 ng, 10 ng, 5 ng, 1 ng), Internal positions DYKDDDDK-tagged fusion protein (9-13: 50 ng, 25 ng, 10 ng, 5 ng, 1 ng respectively), IRDye 800 conjugated Goat anti-Rabbit IgG (H&L) was used as secondary antibody at 0.125  $\mu$ g/ml.




Figure-3 : Western blot analysis of DYKDDDDK Tag Antibody (Clone: 114F12C8) on immunoprecipitation from cell lysates containing DYKDDDDK fusion protein of 1: DYKDDDDK antibody (4  $\mu$ g) + Hela cell lysate containing N-terminal DYKDDDDK tagged fusion protein, 2: Rabbit IgG (4  $\mu$ g)+ Hela cell lysate containing N-terminal DYKDDDDK tagged fusion protein, 3: N-terminal DYKDDDDK tagged fusion protein (Input), 4: DYKDDDDK antibody (4  $\mu$ g) + Hela cell lysate containing C-terminal DYKDDDDK tagged fusion protein, 5: Rabbit IgG (4  $\mu$ g)+ Hela cell lysate containing C-terminal DYKDDDDK tagged fusion protein, 6: C-terminal DYKDDDDK tagged fusion protein (Input) at 0.2  $\mu$ g/ml, IRDye800 conjugated Goat anti-Rabbit IgG (H&L) was used as secondary antibody at 0.125  $\mu$ g/ml.