

## 10-4009: Monoclonal Antibody to IL-17F (Clone: ABM2H17)

|                                |   |
|--------------------------------|---|
| <b>Clonality :</b>             | Monoclonal  |
| <b>Clone Name :</b>            | ABM2H17   |
| <b>Application :</b>           | WB  |
| <b>Reactivity :</b>            | Human   |
| <b>Gene :</b>                  | IL17F   |
| <b>Gene ID :</b>               | 112744  |
| <b>Uniprot ID :</b>            | Q96PD4  |
| <b>Format :</b>                | Purified  |
| <b>Alternative Name :</b>      | Cytokine ML-1, Interleukin-17F  |
| <b>Isotype :</b>               | Mouse IgG1 Kappa  |
| <b>Immunogen Information :</b> | Full length recombinant IL-17F protein was used as the immunogen for this antibody. |

### Product Info

|                            |  |
|----------------------------|--|
| <b>Amount :</b>            | 25 µg / 100 µg   |
| <b>Purification :</b>      | Protein G Chromatography   |
| <b>Content :</b>           | 25 µg in 50 µl/100 µg in 200 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly toxic.               |
| <b>Storage condition :</b> | Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles |

### Application Note

Recommended dilutions: WB: 6-8 µg/ml. However, this need to be optimized based on the research applications.




Figure-1: Western blot analysis of IL-17F. Anti- IL-17F antibody (Clone:ABM2H17) was tested at 6 µg/ml on h Lung lysates.