

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 12-1157: Anti-Cytokeratin 19 (KRT19) (Pancreatic Stem Cell Marker) Recombinant Mouse Monoclonal Antibody (Clone:rKRT19/799)

Clonality: Monoclonal **Clone Name:** rKRT19/799 Application: FACS, IF, WB, IHC

Reactivity: Human Gene: KRT19 Gene ID: 3880 **Uniprot ID:** P08727 Format: Purified

**Alternative Name:** k19; k1cs; Keratin 19 Keratin Type i 40kD; krt19

Isotype: Mouse IgG1, kappa

Immunogen Information: Recombinant full-length human KRT19 protein

## **Description**

This MAb reacts with the rod domain of human cytokeratin-19 (CK19), a polypeptide of 40kDa. Its epitope maps between amino acid 312-335. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epithelium. Anti-CK19 reacts with a wide variety of epithelial malignancies including adenocarcinomas of the colon, stomach, pancreas, biliary tract, liver, and breast. Perhaps the most useful application is the identification of thyroid carcinoma of the papillary type, althoµgh 50%-60% of follicular carcinomas are also labeled. Anti-CK19 is a useful marker for detection of tumor cells in lymph nodes, peripheral blood, bone marrow and breast cancer.

## **Product Info**

Amount: 20 μg / 100 μg **Purification:** Protein A/G

200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Content:

0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody Storage condition:

is stable for 24 months. Non-hazardous.

## **Application Note**

Flow Cytometry (1-2Ă∏Âμg/million cells); Immunofluorescence (1-2Ă∏Âμg/ml); Western Immunohistochemistry (Formalin-fixed) (1-2ÃΠÂμg/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Formalin-fixed, paraffin-embedded Human Colon stained with Cytokeratin 19 Mouse Recombinant Monoclonal Antibody (rKRT19/799).




Figure 2: SDS-PAGE Analysis Purified Cytokeratin 19 Mouse Recombinant Monoclonal Antibody (rKRT19/799).