

12-1213: Anti-Moesin Recombinant Mouse Monoclonal Antibody (Clone:rMSN/492)

Clonality :	Monoclonal
Clone Name :	rMSN/492
Application :	FACS,IF,WB,IHC
Reactivity :	Human, Rat
Gene :	MSN
Gene ID :	4478
Uniprot ID :	P26038
Format :	Purified
Alternative Name :	Membrane-organizing extension spike protein; Moesin/anaplastic lymphoma kinase fusion protein; MSN/ALK fusion
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant full-length human Moesin protein

Description

Recognizes 78kDa moesin protein. Moesin, a member of the talin-4.1 superfamily, is a linking protein of the sub-membranous actin cytoskeleton. It is expressed in variable amounts in cells of different phenotypes such as macrophages, lymphocytes, fibroblastic, endothelial, epithelial, and neuronal cell lines but not in blood cells. The ERM proteins, ezrin, radixin, and moesin are involved in a variety of cellular functions, such as cell adhesion, migration, and the organization of cell surface structures, and are highly homologous, both in protein sequence and in functional activity, with merlin/schwannomin, a neurofibromatosis-2-associated tumor-suppressor protein. Cell lines of epithelial and mesothelial origin contain both moesin and radixin whereas cells of endothelial and lymphoid origin express moesin.

Product Info

Amount :	20 µg / 100 µg
Purification :	Protein A/G
Content :	200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2 \times 10⁵µg/million cells); Immunofluorescence (2-4 \times 10⁵µg/ml); Western Blot (1-2 \times 10⁵µg/ml); Immunohistochemistry (Formalin-fixed) (1-2 \times 10⁵µg/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Figure 1: Formalin-fixed, paraffin-embedded human Placenta stained with Moesin Mouse Recombinant Monoclonal Antibody (rMSN/492).




Figure 2: SDS-PAGE Analysis Purified Moesin Mouse Recombinant Monoclonal Antibody (rMSN/492).