

## 12-1215: Anti-MUC1 / CA15-3 / EMA / CD227 (Epithelial Marker) Recombinant Rabbit Monoclonal Antibody (Clone:MUC1/2278R)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	MUC1/2278R
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	MUC1
<b>Gene ID :</b>	4582
<b>Uniprot ID :</b>	P15941
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Breast carcinoma-associated antigen DF3, CA15-3, Carcinoma-associated mucin Episialin, Epithelial Membrane Antigen, H23AG, KL-6, MAM6, MUC-1, MUC-1/SEC, MUC-1/X, MUC1-alpha, MUC1-beta, MUC1-CT, MUC1-NT, MUC1/ZD, Mucin 1 cell surface associated, Mucin-1 subunit beta, Peanut-reactive urinary mucin, PEM, PEMT, Polymorphic epithelial mucin, PUM, Tumor-associated epithelial membrane antigen
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Human milk-fat globule membranes (HMFGM)

### Description

This MAb reacts with MUC1, a large transmembrane glycoprotein expressed on the ductal surface of normal glandular epithelia. It is used as tracer agent in CA15.3 assays. The extracellular domain of MUC1 largely consists of a highly conserved, O-glycosylated 20 amino acids tandem repeat which can occur 30-100 times per molecule depending on the length of the allele involved. In the vast majority of human carcinomas this protein is up-regulated and poorly glycosylated and appears on the cell surface in a non-polarized fashion. The dominant epitope of this MAb involves both amino acids as well as sÅµgar moieties. Its epitope is destroyed by desialylation i.e. treatment with Neuraminidase.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Purification :</b>	Protein A/G
<b>Content :</b>	200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunohistochemistry (Formalin-fixed) (1-2Åµg/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Figure 1: Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with MUC1 Rabbit Recombinant Monoclonal Antibody (MUC1/2278R).




Figure 2: SDS-PAGE Analysis of Purified MUC1 Rabbit Recombinant Monoclonal Antibody (MUC1/2278R).