

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 12-1223: Anti-Neurofilament (NF-H) (Neuronal Marker) Recombinant Mouse Monoclonal Antibody (Clone:rNF421)

Clone Name: Monoclonal rNF421

Application: IHC,FACS,WB
Reactivity: Human
Gene: NEFH
Gene ID: 4744
Uniprot ID: P12036
Format: Purified

Alternative Name:

NEFH; Neurofilament H; Neurofilament Heavy Polypeptide 200kDa; Neurofilament Triplet H

Protein; NF-H; NF200 **Isotype:** Mouse IgG1, kappa

**Immunogen Information:** Recombinant human neurofilament protein

## **Description**

This MAb reacts with a 200kDa protein, identified as heavy sub-unit of neurofilaments (NF-H). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68kDa (NF-L), 160kDa (NF-M) and 200kDa (NF-H). Anti-neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.

## **Product Info**

Amount :  $20 \mu g / 100 \mu g$ Purification : Protein A/G

Content: 200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA &

0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

**Storage condition :** Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

## **Application Note**

Western Blot (1-2µg/ml); Flow Cytometry (1-2µg/million cells);Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com



Figure 1: Formalin-fixed, paraffin-embedded human Brain stained with Neurofilament Mouse Recombinant Monoclonal Antibody (rNF421).




Figure 2: SDS-PAGE Analysis of Purified Neurofilament Mouse Recombinant Monoclonal Antibody (rNF421).