

## 12-1114: Anti-CD43 (T-Cell Marker) Recombinant Mouse Monoclonal Antibody (Clone:rSPN/1094)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	rSPN/1094
<b>Application :</b>	WB,FACS,IF,IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	SPN
<b>Gene ID :</b>	6693
<b>Uniprot ID :</b>	P16150
<b>Format :</b>	Purified
<b>Alternative Name :</b>	Galactoglycoprotein, GALGP, GPL115, Leukocyte sialoglycoprotein, Leukosialin, LSN, Sialophorin, SPN
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Human full-length recombinant CD43 protein

### Description

It recognizes a cell surface glycoprotein of 95/115/135kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Purification :</b>	Protein A/G
<b>Content :</b>	200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Western Blot (1-2µg/ml); Flow Cytometry (1-2µg/million cells); Immunofluorescence (1-2µg/ml); Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Figure 1: Formalin-fixed, paraffin-embedded human Lymphoma stained with CD43 Mouse Recombinant Monoclonal Antibody (rSPN/1094).




Figure 2: Formalin-fixed, paraffin-embedded human Tonsil stained with CD43 Mouse Recombinant Monoclonal Antibody (rSPN/1094).




Figure 3: SDS-PAGE Analysis Purified CD43 Mouse Recombinant Monoclonal Antibody (rSPN/1094).