

12-1248: Anti-Thyroglobulin (Thyroidal Cell Marker) Recombinant Mouse Monoclonal Antibody (Clone:r2H11)

Clonality :	Monoclonal
Clone Name :	r2H11
Application :	IHC
Reactivity :	Human, Mouse, Rat
Gene :	TG
Gene ID :	7038
Uniprot ID :	P01266
Format :	Purified
Alternative Name :	AITD3, hTG, TDH3, Tg, Tgn
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Human thyroid follicular cells

Description

MAb r2H11 reacts with a partially defined epitope of human thyroglobulin. This epitope is different from the epitope recognized by MAb 6E1. Thyroglobulin is a 660kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo conformational maturation in the endoplasmic reticulum. The vast majority of follicular carcinomas of the thyroid will give positive immunoreactivity for anti-thyroglobulin even though sometimes only focally. Poorly differentiated carcinomas of the thyroid are frequently anti-thyroglobulin negative. Adenocarcinomas of other-than-thyroid origin do not react with this antibody. This antibody is useful in identification of thyroid carcinoma of the papillary and follicular types. Presence of thyroglobulin in metastatic lesions establishes the thyroid origin of tumor. Anti-thyroglobulin, combined with anti-calcitonin, can identify medullary carcinomas of the thyroid. Furthermore, anti-thyroglobulin, combined with anti-TTF1, can be a reliable marker to differentiate between primary thyroid and lung neoplasms.

Product Info

Amount :	20 µg / 100 µg
Purification :	Protein A/G
Content :	200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (0.1-0.2 µg/ml; 30 min at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Figure 1: Formalin-fixed, paraffin-embedded human Thyroid stained with Thyroglobulin Mouse Recombinant Monoclonal Antibody (r2H11).




Figure 2: SDS-PAGE Analysis of Purified Thyroglobulin Mouse Recombinant Monoclonal Antibody (r2H11).