

12-1216: Anti-MUC16 / CA125 (Ovarian Carcinoma Marker) Recombinant Rabbit Monoclonal Antibody (Clone:OCA125/2349R)

Clonality :	Monoclonal
Clone Name :	OCA125/2349R
Application :	IHC
Reactivity :	Human
Gene :	CA125
Gene ID :	94025
Uniprot ID :	Q8WXI7
Format :	Purified
Alternative Name :	CA125; Cancer antigen 125; MUC16; Mucin16; Ovarian cancer-related tumor marker CA125; Ovarian carcinoma antigen CA125
Isotype :	Rabbit IgG
Immunogen Information :	Recombinant full-length human MUC16 protein

Description

The mucins are a family of highly glycosylated, secreted proteins with a basic structure consisting of a variable number of tandem repeats (VNTRs). Membrane-associated and secretory Mucins are high molecular weight glycoproteins of the glycocalyx (polysaccharide biofilm) that protects mucosal epithelium from particulate matter and microorganisms. Epithelial Mucins are large, secreted and cell surface glycoproteins crucial for adhesion modulation, signaling and epithelial cell protection. The number of repeats is highly polymorphic and varies among different alleles. The Mucin family consists of Mucins 1-4, Mucin 5 (AC and B), Mucins 6-8, Mucins 11-13 and Mucins 15-17. The Mucin 16 protein (also commonly referred to as CA125), encoded for by the gene MUC16, is a very high molecular weight tumor antigen consisting of three domains: a carboxy terminal domain, an extracellular domain and an amino terminal domain. Mucin 16, an ovarian cancer-associated antigen, is used as a marker to monitor the progress of epithelial ovarian cancer. It is a hydrophilic membrane-associated protein that may be involved in vitamin A functions.

Product Info

Amount :	20 µg / 100 µg
Purification :	Protein A/G
Content :	200µg/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Figure 1: Formalin-fixed, paraffin-embedded human Ovarian Carcinoma stained with MUC16 Rabbit Recombinant Monoclonal Antibody (OCA125/2349R).




Figure 2: Formalin-fixed, paraffin-embedded human Endometrial Carcinoma stained with MUC16 Rabbit Recombinant Monoclonal Antibody (OCA125/2349R).




Figure 3: SDS-PAGE Analysis of Purified MUC16 Rabbit Recombinant Monoclonal Antibody (OCA125/2349R).