

## 30-1964: FITC Conjugated, Anti-alpha-tubulin Monoclonal Antibody (Clone:TU-01)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	TU-01
<b>Application :</b>	WB
<b>Reactivity :</b>	Broad species reactivity
<b>Conjugate :</b>	FITC
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Fraction of tubulin purified from porcine brain by two cycles of polymerization - depolymerization.

### Description

The microtubules are intracellular dynamic polymers made up of evolutionarily conserved polymorphic alpha/beta-tubulin heterodimers and a large number of microtubule-associated proteins (MAPs). The microtubules consist of 13 protofilaments and have an outer diameter 25 nm. Microtubules have their intrinsic polarity; highly dynamic plus ends and less dynamic minus ends. Microtubules are required for vital processes in eukaryotic cells including mitosis, meiosis, maintenance of cell shape and intracellular transport. Microtubules are also necessary for movement of cells by means of flagella and cilia. In mammalian tissue culture cells microtubules have their minus ends anchored in microtubule organizing centers (MTOCs). The GTP (guanosinotriphosphate) molecule is an essential for tubulin heterodimer to associate with other heterodimers to form microtubule. In vivo, microtubule dynamics vary considerably. Microtubule polymerization is reversible and a populations of microtubules in cells are on their minus ends either growing or shortening - this phenomenon is called dynamic instability of microtubules. On a practical level, microtubules can easily be stabilized by the addition of non-hydrolysable analogues of GTP (eg. GMPPCP) or more commonly by anti-cancer drugs such as Taxol. Taxol stabilizes microtubules at room temperature for many hours. Using limited proteolysis by enzymes both tubulin subunits can be divided into N-terminal and C-terminal structural domains. The alpha-tubulin (relative molecular weight around 50 kDa) is globular protein that exists in cells as part of soluble alpha/beta-tubulin dimer or it is polymerized into microtubules. In different species it is coded by multiple tubulin genes that form tubulin classes (in human 6 genes). Expressed tubulin genes are named tubulin isotypes. Some of the tubulin isotypes are expressed ubiquitously, while some have more restricted tissue expression. Alpha-tubulin is also subject of numerous post-translational modifications. Tubulin isotypes and their posttranslational modifications are responsible for multiple tubulin charge variants - tubulin isoforms. Heterogeneity of alpha-tubulin is concentrated in C-terminal structural domain.

### Product Info

<b>Amount :</b>	0.1 mg
<b>Storage condition :</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Immunofluorescence staining of 3T3 mouse embryonal fibroblast cell line using anti-alpha-tubulin (TU-01; green) and anti-Vimentin (VI-01; ). Nucleus is stained with DAPI (blue).




Figure 2: Immunofluorescence staining of HeLa human cervix carcinoma cell line using anti-alpha-tubulin (TU-01; red). Nucleus is stained with DAPI (blue).




Figure 3: Immunofluorescence staining of 3T3 mouse embryonal fibroblast cell line using anti-alpha-tubulin (TU-01; green). Nucleus is stained with DAPI (blue).




Figure 4: Use of anti-alpha-tubulin antibody TU-01 as a loading control (A) in a Western blotting experiment revealing the staining pattern of various cell lysates by a newly developed monoclonal antibody (B).




Figure 5: Immunoprecipitation of alpha-tubulin from HeLa and RAJI cell lysate by antibody TU-16 and its detection by antibody TU-01. IgM heavy chain (76-92 kDa) and IgM light chain (25-30 kDa) indicated. Mr of alpha tubulin is around 50 kDa. L = lysate IPr = immunoprecipitate (reducing conditions) IPn = immunoprecipitate (non-reducing conditions)