

30-1965: FITC Conjugated, Anti-betaIII-tubulin Monoclonal Antibody (Clone:TU-20)

Clonality :	Monoclonal
Clone Name :	TU-20
Application :	FACS
Reactivity :	Broad species reactivity
Conjugate :	FITC
Isotype :	Mouse IgG1
Immunogen Information :	Peptide (C) 441-448 coupled to maleimide-activated keyhole limpet hemocyanin via cysteine added to the N-terminus of the neuron-specific peptide.

Description

The betaIII-tubulin isoform is present dominantly in cells of neuronal origin and it is one of the earliest markers of neuronal differentiation. It is regarded as a specific probe for the cells of neuronal origin as well as for the tumours originating from these cells. The betaIII-tubulin is most abundant in cells of neuronal origin, but was also detected in Sertoli cells of the testis and transiently in non-neuronal embryonic tissues.

Product Info

Amount :	0.1 mg
Storage condition :	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Immunofluorescence staining (mouse neuroblastoma cells) Figure 1: Immunofluorescence staining of Neuro2a mouse neuroblastoma cell line using anti-betaIII-tubulin (TU-20; green; 3 µg/ml). Nuclei were stained with DAPI (blue).




Figure 2: Immunofluorescence staining of P-19 mouse embryonal carcinoma cell line stimulated to neuronal differentiation by retinoic acid. 2A - Microtubules decorated with neuron-specific anti-betaIII-tubulin (TU-20; red). 2B - Merged image of co-staining with anti-beta-tubulin (TU-06; green;). Superposition of red and green colours provided yellow staining. Nuclei were stained with DNA-binding dye (blue).