

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-1397: Anti-Bim Monoclonal Antibody (Clone: Ham151-149) (Discontinued)

Clonality: Monoclonal
Clone Name: Ham151-149

Application: FACS
Reactivity: Mouse
Gene: Bcl2|11
Gene ID: 12125
Uniprot ID: 054918
Format: Purified

Alternative Name: Bcl2-interacting mediator of cell death

**Isotype:** Hamster IgG1

Immunogen Information: Mouse BimL (w/o transmembrane region) + Bcl-xL (w/o transmembrane region)-6-his tag,

purified from Hi5 cells

## **Description**

Bim (Bcl2-interacting mediator) is a pro-apoptotic protein of BH3 domain-only subgroup of the Bcl2 family. It has important roles in initiation of apoptosis in response to many death stimuli. Bim is an important regulator of B and T cell negative selection and is also an essential regulator of T cell apoptosis during termination of an immune response. Bim is constitutively expressed in many cell types but it is maintained in an inactive form through binding to the microtubule-associated dynein motor complex.

## **Product Info**

Amount: 0.1 mg

**Purification :** Purified by protein-A affinity chromatography

**Storage condition :** Store at 2-8°C. Do not freeze.

## **Application Note**

Flow Cytometry Recommended dilution:10  $\mu g$  / ml

Application note: Cells permeabilized with 0.03% saponin

**Immunoprecipitation** Recommended dilution: 1-2  $\mu$ g /100-500  $\mu$ g of protein in 1 ml lysate **Western Blotting** Recommended dilution: 1-2  $\mu$ g / ml Application note: reducing conditions

Immunocytochemistry Application note: Permeabilized cells