∗ abeomics

30-2417: Anti-Blood Group ABH culture supernatant Monoclonal Antibody (Clone:HE-10)

Clonality :	Monoclonal
Clone Name :	HE-10
Application :	IHC,FACS
Reactivity :	Human
Isotype :	Mouse IgM
Immunogen Information	Mixture of erythrocytes of group A1 and glycoprotein fraction isolated from saliva of secretors with blood group A.

Description

The mouse monoclonal antibody HE-10 agglutinates erythrocytes of group A, and is excellent as a tumour marker in patients of blood group B and 0. It does not agglutinate erythrocytes of group B and 0. Study with specific oligosaccharides showed that the antibody HE-10 reacts with A and H antigens with chain types 3 and 4 and it does not react with A disaccharide, A trisaccharide, A type 1, A type 2, ALeb. The antibody HE-10 does not react with normal tissue sections of donors with blood group B and 0 but it reacts specifically with malignant tissues. The mouse monoclonal antibody HE-10 agglutinates erythrocytes of group A, and is excellent as a tumour marker in patients of blood group B and 0. It does not agglutinate erythrocytes of group B and 0. Study with specific oligosaccharides showed that the antibody HE-10 reacts with A and H antigens with chain types 3 and 4 and it does not react with A disaccharide, A trisaccharide, A type 1, A type 2, ALeb. The antibody HE-10 reacts with A and H antigens with chain types 3 and 4 and it does not react with A disaccharide, A trisaccharide, A type 1, A type 2, ALeb. The antibody HE-10 does not reacts specifically with malignant tissue sections of donors with blood group B and 0 but it reacts specifies and 4 and it does not react with A disaccharide, A trisaccharide, A type 1, A type 2, ALeb. The antibody HE-10 does not react with normal tissue sections of donors with blood group B and 0 but it reacts specifically with malignant tissues.

The mouse monoclonal antibody HE-10 agglutinates erythrocytes of group A, and is excellent as a tumour marker in patients of blood group B and O. It does not agglutinate erythrocytes of group B and O. Study with specific oligosaccharides showed that the antibody HE-10 reacts with A and H antigens with chain types 3 and 4 and it does not react with A disaccharide, A trisaccharide, A type 1, A type 2, ALeb. The antibody HE-10 does not react with normal tissue sections of donors with blood group B and O but it reacts specifically with malignant tissues.

Product Info

Amount :1.0 mlStorage condition :Store at 2-8°C. Do not freeze.

₩ abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com

