

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-1118: Anti-CD14 / LPS-receptor Monoclonal Antibody (Clone:MEM-18)

Clonality: Monoclonal
Clone Name: MEM-18
Application: FACS

**Reactivity:** Non-Human Primates

Gene : CD14
Gene ID : 929
Uniprot ID : P08571
Format : Purified
Alternative Name : CD14
Isotype : Mouse IgG1

Immunogen Information: A crude mixture of human urinary proteins precipitated by ammonium sulphate from the urine

of a patient suffering from proteinuria.

## **Description**

The antibody MEM-18 reacts with CD14, a 53-55 kDa GPI (glycosylphosphatidylinositol)-linked extracellular membrane glycoprotein expressed on monocytes, macrophages and weakly on granulocytes; also expressed by most tissue macrophages. In human, the epitope recognized by MEM-18 is located between amino acids 57-64.

The antibody MEM-18 reacts with CD14, a 53-55 kDa GPI (glycosylphosphatidylinositol)-linked extracellular membrane glycoprotein expressed on monocytes, macrophages and weakly on granulocytes; also expressed by most tissue macrophages. In human, the epitope recognized by MEM-18 is located between amino acids 57-64.

# **Product Info**

Amount: 0.1 mg

**Purification:** Purified by protein-A affinity chromatography

**Storage condition :** Store at 2-8°C. Do not freeze.

### **Application Note**

#### Flow Cytometry Recommended dilution:4 µg/ml

Immunoprecipitation Western Blotting ELISA The antibody MEM-18 has been tested as the detection antibody in a sandwich ELISA for analysis of human CD14 in combination with antibody B-A8 . **Functional Application** The antibody MEM-18 completely blocks binding of fluorescein (FITC) - labeled bacterial LPS to the monocyte surface and it also blocks the binding of CD14 to the extracellular TLR2 domain.