

30-1005: Anti-CD16 / FcgammaRIII Monoclonal Antibody (Clone:MEM-154)-Azide free

Clonality :	Monoclonal
Clone Name :	MEM-154
Application :	FACS
Reactivity :	Human
Gene :	FCGR3A
Gene ID :	2214
Uniprot ID :	P08637
Alternative Name :	FCGR3A,CD16A,FCG3,FCGR3,IGFR3
Isotype :	Mouse IgG1
Immunogen Information : Human granulocytes	

Description

CD16 (FcgammaRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcgammaRIII is expressed in two forms - FcgammaRIII-A and -B. FcgammaRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcepsilonRI-gamma subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcgammaRIII-A is associated, moreover, with FcepsilonRI-beta subunit. Besides IgG, FcgammaRIII-A can be triggered also by oligomeric IgE. FcgammaRIII-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

Product Info

Amount :	0.1 mg
Purification :	Purified by protein-A affinity chromatography
Storage condition :	Store at 2-8°C. Do not freeze.

Application Note

Flow Cytometry Recommended dilution: 5-10 µg/ml

Positive control: PBL (peripheral blood lymphocytes)

Application note: The antibody MEM-154 does not react with CD16a present on NK cells in many subjects. Immunoprecipitation Western Blotting Application note: Non-reducing conditions. Functional Application The antibody MEM-154 blocks binding of human IgG to FcgammaRIII.




Figure 1: Flow cytometry surface staining pattern of human peripheral blood stained using anti-human CD16 (MEM-154) purified antibody (azide free, concentration in sample 2 μg/ml) GAM APC.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

₩ abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Figure 2: Separation of human neutrophil granulocytes (red-filled) from CD16 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD16 (MEM-154) purified antibody (azide free, concentration in sample 2 \hat{I}_{4} g/ml) GAM APC.