

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-1120: Anti-CD16 / FcgammaRIII Monoclonal Antibody (Clone:MEM-154)

Clonality: Monoclonal **Clone Name:** MEM-154 Application: FACS Reactivity: Human Gene: FCGR3A Gene ID: 2214 **Uniprot ID:** P08637 **Purified** Format:

Alternative Name: FCGR3A,CD16A,FCG3,FCGR3,IGFR3

Isotype: Mouse IgG1

Immunogen Information: Human granulocytes

Description

CD16 (FcgammaRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcgammaRIII is expressed in two forms - FcgammaRIII-A and -B. FcgammaRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcepsilonRI-gamma subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcgammaRIII-A is associated, moreover, with FcepsilonRI-beta subunit. Besides IgG, FcgammaRIII-A can be triggered also by oligomeric IgE. FcgammaRIII-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

Product Info

Amount: 0.1 mg

Purification: Purified by protein-A affinity chromatography

Storage condition : Store at 2-8°C. Do not freeze.

Application Note

Flow Cytometry Recommended dilution: 5-10 μg/ml Positive control: PBL (peripheral blood lymphocytes)

Application note: The antibody MEM-154 does not react with CD16a present on NK cells in many subjects. **Immunoprecipitation Western Blotting** Application note: Non-reducing conditions. **Functional Application** The antibody MEM-154 blocks binding of human IgG to FcgammaRIII.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 1: Flow cytometry surface staining pattern of human peripheral blood stained using anti-human CD16 (MEM-154) purified antibody (concentration in sample 2 $\hat{1}\frac{1}{4}$ g/ml) GAM APC.




Figure 2: Separation of FasL transfected L5178Y cells stained using anti-human CD178 (NOK-1) purified antibody (concentration in sample 9 $\hat{1}\frac{1}{4}$ g/ml, GAM APC, red-filled) from FasL transfected L5178Y cells unstained by primary antibody (GAM APC, black-dashed) in flow cytometry analysis (surface staining).