

**30-1258: Anti-CD16 / Fcγ<sub>3</sub> Monoclonal Antibody (Clone:MEM-168)**

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	MEM-168
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human, Non-Human Primates, Pig
<b>Gene :</b>	FCGR3A
<b>Gene ID :</b>	2214
<b>Uniprot ID :</b>	P08637
<b>Format :</b>	Purified
<b>Alternative Name :</b>	FCGR3A,CD16A,FCG3,FCGR3,IGFR3
<b>Isotype :</b>	Mouse IgM
<b>Immunogen Information :</b>	Human granulocytes

**Description**

CD16 (Fcγ<sub>3</sub>) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human Fcγ<sub>3</sub> is expressed in two forms - Fcγ<sub>3</sub>-A and -B. Fcγ<sub>3</sub>-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcεR1-gamma subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell Fcγ<sub>3</sub>-A is associated, moreover, with FcεR1-beta subunit. Besides IgG, Fcγ<sub>3</sub>-A can be triggered also by oligomeric IgE. Fcγ<sub>3</sub>-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

**Product Info**

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified by precipitation and chromatography
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.

**Application Note**

**Flow Cytometry** *Recommended dilution:* 1-4  $\mu$ g/ml