

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-1599: Anti-CD18 / Integrin beta2 subunit Monoclonal Antibody (Clone:MEM-48)-Low Endotoxin

Clonality: Monoclonal Clone Name: MEM-48

**Application :** FACS, IP, WB, IHC, IHC-Fr, Functional Assay

Reactivity: Human
Gene: ITGB2
Gene ID: 3689
Uniprot ID: P05107

Format: Low Endotoxin

Alternative Name: ITGB2,CD18,MFI7

Isotype: Mouse IgG1

Immunogen Information: Leukocytes of a patient suffering from a LGL-type leukemia.

## **Description**

CD18, integrin beta2 subunit, forms heterodimers with four types of CD11 molecule to constitute leukocyte (beta2) integrins: alphaLbeta2 (CD11a/CD18, LFA-1), alphaMbeta2 (CD11b/CD18, Mac-1, CR3), alphaXbeta2 (CD11c/CD18) and alphaDbeta2 (CD11d/CD18). In most cases, the response mediated by the integrin is a composite of the functions of its individual subunits. These integrins are essential for proper leukocyte migration, mediating intercellular contacts. Absence of CD18 leads to leukocyte adhesion deficiency-1; severe reduction of CD18 expression leads to the development of a psoriasiform skin disease. CD18 is also a target of Mannheimia (Pasteurella) haemolytica leukotoxin and is sufficient to mediate leukotoxin-mediated cytolysis.

## **Product Info**

Amount: 0.1 mg

**Purification :** Purified by protein-A affinity chromatography

**Storage condition :** Store at 2-8°C. Do not freeze.

## **Application Note**

Western Blotting Recommended dilution: 2 ̸µg/ml Positive control: Kg-1a human leukemia cell lysate

Immunohistochemistry Recommended dilution: 10 ̸µg/ml

Positive tissue: spleen, microglia




Figure 1:Western Blotting analysis (non-reducing conditions) of whole cell lysate using anti-human CD18 (MEM-48). Lane 1: KG-1a human acute myeloid leukemia cell line Lane 2: HPB-ALL human peripheral blood T cell leukemia cell line



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure 2: Immunohistochemistry staining of human spleen (paraffin sections) using anti-CD18 (MEM-48).