

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 30-2168: Anti-CD29 / Integrin beta1 chain Monoclonal Antibody (Clone:MEM-101A)-PE Conjugated

Clonality: Monoclonal
Clone Name: MEM-101A
Application: FACS, IP, WB
Reactivity: Human, Pig, Dog

 Conjugate :
 PE

 Gene :
 ITGB1

 Gene ID :
 3688

 Uniprot ID :
 P05556

**Alternative Name:** ITGB1,FNRB,MDF2,MSK12

**Isotype:** Mouse IgG1

Immunogen Information: Raji Burkitt's lymphoma cell line

## **Description**

CD29 (beta1 integrin subunit, GPIIa) forms non-covalently linked heterodimers with at least 6 different alpha chains (alpha1-alpha6, CD49a-f) determining the binding properties of beta1 (VLA) integrins. These integrins mediate cell adhesion to collagen, fibronectin, laminin and other extracellular matrix (ECM) components. This interaction hinders cell death, whereas disruption of anchorage to ECM leads to apoptosis. Decreased expression of most beta1 integrins correlates with acquiring multidrug resistance of tumour cells during selection in presence of antitumour drug. In platelets, translocation of intracellular pool of beta1 integrins to the plasma membrane following thrombin stimulation. These integrins are also upregulated in leukocytes during emigration and extravascular migration and appear to be critically involved in regulating the immune cell trafficking from blood to tissue, as well as in regulating tissue damage and disease symptoms related to inflammatory bowel disease. Through a beta1 integrin-dependent mechanism, fibronectin and type I collagen enhance cytokine secretion of human airway smooth muscle in response to IL-1beta.

## **Product Info**

Amount: 100 tests

**Storage condition:** Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Surface staining of human peripheral blood with anti-human CD29 (MEM-101A) PE.