

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

30-2240: Anti-CD30 / Ki-1 Monoclonal Antibody (Clone:MEM-268)-PE Conjugated

Clonality: Monoclonal
Clone Name: MEM-268
Application: FACS
Reactivity: Human
Conjugate: PE
Gene: TNFRSF8
Gene ID: 943

Alternative Name: TNFRSF8,CD30,D1S166E

P28908

Isotype: Mouse IgG

Immunogen Information: Expression vector containing CD30 cDNA (booster suspension of THP-1 cell line)

Description

Uniprot ID:

CD30 is a type I transmembrane glycoprotein of the TNF receptor superfamily. CD30 was originally identified as a cell surface antigen of Hodgkins and Reed-Sternberg cells using monoclonal antibody Ki-1. The ligand for CD30 is CD30L (CD153). The binding of CD30 to CD30L mediates pleiotropic effects including cell proliferation, activation, differentiation, and apoptotic cell death. CD30 has a critical role in the pathophysiology of Hodgkin's disease and other CD30+ lymphomas. CD30 acts as a costimulatory molecule in thymic negative selection. In addition to its expression on Hodgkin's and Reed-Sternberg cells, CD30 is also found in some non-Hodgkin's lymphomas (including Burkitt's lymphomas), virus-infected T and B cells, and on normal T and B cells after activation. In T cells, CD30 expression is present on a subset of T cells that produce Th2-type cytokines and on CD4+/CD8+ thymocytes that co-express CD45RO and the IL4 receptor. Soluble form of CD30 (sCD30) serves as a marker reflecting Th2 immune response.

Product Info

Amount: 100 tests

Content: Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide **Storage condition:** Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.

Application Note

Flow cytometry: The reagent is designed for analysis of human blood cells using 20 μ l reagent / 100 μ l of whole blood or 106 cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.

Flow cytometry: The reagent is designed for analysis of human blood cells using 20 μ l reagent / 100 μ l of whole blood or 106 cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.