

30-1109: Anti-CD4 Monoclonal Antibody (Clone:MEM-115)

Clonality :	Monoclonal
Clone Name :	MEM-115
Application :	IP
Reactivity :	Human
Gene :	CD4
Gene ID :	920
Uniprot ID :	P01730
Format :	Purified
Alternative Name :	CD4
Isotype :	Mouse IgG2a
Immunogen Information :	Human thymocytes and T lymphocytes.

Description

CD4 is a single chain transmembrane glycoprotein and belongs to immunoglobulin supergene family. In extracellular region there are 4 immunoglobulin-like domains (1 Ig-like V-type and 3 Ig-like C2-type). Transmembrane region forms 25 aa, cytoplasmic tail consists of 38 aa. Domains 1,2 and 4 are stabilized by disulfide bonds. The intracellular domain of CD4 is associated with p56Lck, a Src-like protein tyrosine kinase. It was described that CD4 segregates into specific detergent-resistant T-cell membrane microdomains. Extracellular ligands: MHC class II molecules (binds to CDR2-like region in CD4 domain 1); HIV envelope protein gp120 (binds to CDR2-like region in CD4 domain 1); IL-16 (binds to CD4 domain 3), Human seminal plasma glycoprotein gp17 (binds to CD4 domain 1), L-selectin Intracellular ligands: p56LckCD4 is a co-receptor involved in immune response (co-receptor activity in binding to MHC class II molecules) and HIV infection (human immunodeficiency virus; CD4 is primary receptor for HIV-1 surface glycoprotein gp120). CD4 regulates T-cell activation, T/B-cell adhesion, T-cell differentiation, T-cell selection and signal transduction. Defects in antigen presentation (MHC class II) cause dysfunction of CD4+ T-cells and their almost complete absence in patients blood, tissue and organs (SCID immunodeficiency).

Product Info

Amount :	0.1 mg
Purification :	Purified by protein-A affinity chromatography
Storage condition :	Store at 2-8°C. Do not freeze.

Application Note

Immunoprecipitation Recommended dilution:3 µg/ml




Figure 1: Immunoprecipitation of human CD4 from the lysate of T cells isolated from fresh buffy coats. Western blot was immunostained by anti-human CD4 (MEM-241;). Lane 1: original lysate of T cells. Lane 2: immunoprecipitate by negative control antibody. Lane 3: immunoprecipitate by anti-human CD4 (MEM-115)