

## 30-1990: Anti-CD4 Monoclonal Antibody (Clone:MEM-241)-FITC Conjugated

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	MEM-241
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Conjugate :</b>	FITC
<b>Gene :</b>	CD4
<b>Gene ID :</b>	920
<b>Uniprot ID :</b>	P01730
<b>Alternative Name :</b>	CD4
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	2 N-terminal domains of human CD4 fused to human IgG1 Fc

### Description

CD4 (T4) is a single chain transmembrane glycoprotein and belongs to immunoglobulin supergene family. In extracellular region there are 4 immunoglobulin-like domains (1 Ig-like V-type and 3 Ig-like C2-type). Transmembrane region forms 25 aa, cytoplasmic tail consists of 38 aa. Domains 1,2 and 4 are stabilized by disulfide bonds. The intracellular domain of CD4 is associated with p56Lck, a Src-like protein tyrosine kinase. It was described that CD4 segregates into specific detergent-resistant T-cell membrane microdomains. Extracellular ligands: MHC class II molecules (binds to CDR2-like region in CD4 domain 1); HIV envelope protein gp120 (binds to CDR2-like region in CD4 domain 1); IL-16 (binds to CD4 domain 3), Human seminal plasma glycoprotein gp17 (binds to CD4 domain 1), L-selectin. Intracellular ligands: p56LckCD4 is a co-receptor involved in immune response (co-receptor activity in binding to MHC class II molecules) and HIV infection (human immunodeficiency virus; CD4 is primary receptor for HIV-1 surface glycoprotein gp120). CD4 regulates T-cell activation, T/B-cell adhesion, T-cell differentiation, T-cell selection and signal transduction. Defects in antigen presentation (MHC class II) cause dysfunction of CD4+ T-cells and their almost complete absence in patients blood, tissue and organs (SCID immunodeficiency).

### Product Info

<b>Amount :</b>	100 tests
<b>Storage condition :</b>	Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light.




Figure 1: Surface staining of human peripheral blood cells with anti-human CD4 (MEM-241) FITC.




Figure 2: Separation of human CD4 positive lymphocytes (red-filled) from human CD4 negative neutrophil granulocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD4 (MEM-241) FITC (20  $\mu$ l reagent / 100  $\mu$ l of peripheral whole blood).