

30-1483: Anti-CD4 Monoclonal Antibody (Clone:OX-35)

| | |
|--------------------------------|----------------------------|
| Clonality : | Monoclonal |
| Clone Name : | OX-35 |
| Application : | FACS, IHC, ICC |
| Reactivity : | Rat |
| Gene : | Cd4 |
| Gene ID : | 24932 |
| Uniprot ID : | P05540 |
| Format : | Purified |
| Alternative Name : | Cd4 |
| Isotype : | Mouse IgG2a |
| Immunogen Information : | MLR generated rat Th cells |

Description

CD4 is a single chain transmembrane glycoprotein of immunoglobulin supergene family. In its extracellular region there are 4 immunoglobulin-like domains (1 Ig-like V-type and 3 Ig-like C2-type). The intracellular region of CD4 associates with p56Lck, a Src-like protein tyrosine kinase. It was described that CD4 segregates into specific detergent-resistant T-cell membrane microdomains. CD4 binds to MHC class II molecules (by CDR2-like region in CD4 domain 1), HIV envelope protein gp120 (by CDR2-like region in CD4 domain 1) and other ligands, such as IL-16 (by to CD4 domain 3) or L-selectin. CD4 is a co-receptor involved in immune response (co-receptor activity in binding to MHC class II molecules) and HIV infection. CD4 regulates T-cell activation, T/B-cell adhesion, T-cell differentiation, T-cell selection and signal transduction. Defects in antigen presentation (MHC class II) cause dysfunction of CD4+ T-cells and their almost complete absence in patients blood, tissue and organs (SCID immunodeficiency).

Product Info

| | |
|----------------------------|---|
| Amount : | 0.1 mg |
| Purification : | Purified by protein-A affinity chromatography |
| Storage condition : | Store at 2-8°C. Do not freeze. |