

### 30-1134: Anti-CD54 / ICAM-1 Monoclonal Antibody (Clone:MEM-112)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	MEM-112
<b>Application :</b>	FACS
<b>Reactivity :</b>	Human
<b>Gene :</b>	ICAM1
<b>Gene ID :</b>	3383
<b>Uniprot ID :</b>	P05362
<b>Format :</b>	Purified
<b>Alternative Name :</b>	ICAM1
<b>Isotype :</b>	Mouse IgG1
<b>Immunogen Information :</b>	Raji Burkitt's lymphoma cell line Raji

#### Description

CD54 (ICAM-1) is a 90 kD member of the C2 subset of immunoglobulin superfamily. It is a transmembrane molecule with 7 potential N-glycosylated sites, expressed on resting monocytes and endothelial cells and can be upregulated on many other cells, e.g. with lymphokines, on B- and T-lymphocytes, thymocytes, dendritic cells and also on keratinocytes, chondrocytes, as well as epithelial cells. CD54 mediates cell adhesion by binding to integrins CD11a/CD18 (LFA-1) and to CD11b/CD18 (Mac-1). The interaction of CD54 with LFA-1 enhances antigen-specific T-cell activation.

#### Product Info

<b>Amount :</b>	0.1 mg
<b>Purification :</b>	Purified by protein-A affinity chromatography
<b>Storage condition :</b>	Store at 2-8°C. Do not freeze.

#### Application Note

Flow cytometry: Recommended dilution: 0.5-4 µg/ml.